

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/255720624>

The Harlequin Mantella *Mantella cowanii* in Antoetra Region, High Plateau of Madagascar: New Steps in Conservation

Article · August 2013

CITATIONS

2

READS

286

6 authors, including:



Nirhy Rabibisoa

Universite de Mahajanga

32 PUBLICATIONS 1,271 CITATIONS

SEE PROFILE



Harison Randrianasolo

Conservation International

18 PUBLICATIONS 468 CITATIONS

SEE PROFILE



Falitiana Rabemananjara

Langaha Association

61 PUBLICATIONS 897 CITATIONS

SEE PROFILE



Franco Andreone

Museo Regionale di Scienze Naturali

474 PUBLICATIONS 6,752 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



The invasive Asian common toad (*Duttaphrynus melanostictus*) in Madagascar [View project](#)



IUCN Global Amphibian Assessment 2 [View project](#)

Issue number 107 (July 2013)

ISSN: 1026-0269

eISSN: 1817-3934

FrogLog

www.amphibians.org

Volume 21, number 3

Promoting Conservation, Research and
Education for the World's Amphibians

The Global *Bd*
Mapping Project

Awareness and
Citizen-based
Initiatives

Sabin Award

Recent
Publications

And Much More.



Boophis pyrrhus. Photo by: Jörn Köhler



Amphibian
Survival Alliance
Website



Urban Amphibians
and the Challenges
They Face

The Harlequin Mantella *Mantella cowanii* in Antoetra Region, High Plateau of Madagascar: New Steps in Conservation

By ¹Nirhy H. Rabibisoa, ¹Mirana Anjeriniaina, ²Voahanginirina Rakotoniaina, ¹Harison Randrianasolo, ¹Falitiana E. C. Rabemananjara & ²Franco Andreone



Fig. 1. Fohisokina (or Vohisokina) Massif, selected habitat of *Mantella cowanii* in the Antoetra region, with view of the artificial lake where carp are bred for local consumption.

Since 2008, the conservation program of the Harlequin mantella (*Mantella cowanii*) in Fohisokina site (or Vohisokina), located in the Antoetra Region (Fig. 1), is moving on and scheduled activities are currently shared with the local association “Fohisokina Miaro ny Sahonamena” (= “Fohisokina protects the red frog,” abbreviated FOMISAME) to get a better protection of the site following the species action plan (1). Positive results are now evident and actions need to be continued and reinforced.

Population monitoring

One of us (M. A.) is carrying out her PhD thesis at the University of Antananarivo. The study concerns several aspects the ecology, distribution and population abundance of *M. cowanii* in Fohisokina. The used technique consists of catching the encountered individuals by quadrat sampling. The survey is usually carried out in the early morning (h. 05:00-09:00) before the temperature rises too much, and when frogs are more active. Each individual is photographed and recognized by checking its belly pattern (Fig. 2). In 2012 the total number of captured individuals of *M. cowanii* was 191 within a 2,500 m² surface. The consequent estimation is around 750 individuals per hectare. This number suggests a positive evolution of the population when compared to data at the beginning of the conservation activities in 2008, when only 40 individuals were counted. Furthermore, regular ecological citizen-based



Fig. 3. School construction under progress.



Fig. 2. Belly pattern of *Mantella cowanii*, used to recognize the individuals.

surveys carried out by trained guides of FOMISAME were done every day in the warm-rainy season (October-April). This activity also supported the program on the patrol against illicit activities (i.e., gold mining extraction) within Fohisokina. Twice a year a session of the chytrid monitoring action is also being done (2).

Communication and public awareness

A booklet narrating the history of Fohisokina was published in 2011 by ASG and FOMISAME with the financial help of Conservation International Madagascar (3). It has been developed to increase the knowledge and the importance of the Fohisokina area to people. This handbook has been distributed to schools around Fohisokina and now it is used as a manual (book) in the Ivato municipality. In addition, footage was realized in 2012 with the help of CI, reporting the chronicles of the conservation program on *M. cowanii* with the participation of the local population in Fohisokina. A school building is currently under construction in the Ampadirana village around Fohisokina (Fig. 3). This construction is funded by the NGO, Man and The Environment.

Plant nursery and water quality

Fifteen hundred essential oil plants belonging to the species *Ravensara madagascariensis* were planted around the core site. The aim of this plantation was to improve the livelihood of local populations and mitigate the impact of climate change caused by bushfires. Moreover, the native

Haroungana madagascariensis was planted and cultivated in the buffer zone, close to the core of *M. cowanii* site.

The water sources from Fohisokina supplies an artificial 1-hectare fishpond (carp) realized in 2009 (Fig. 5). This was done as a measure of compensation and to generate revenue. In 2013 the local community has begun to profit from the first production of carp through conservation of the water services.

The water quality is another aspect related to *M. cowanii* conservation. A study to help better understand the hydrographic watershed quality in Fohisokina was completed in February 2013 by a student from Antananarivo University. The objective was to get physico-chemical indications of the pollution level according to potability and drinking water legislation of Madagascar. Fohisokina has five permanent sources for which the main physico-chemical factors were analyzed, as well as quantitative hydrological factors (i.e., temperature, depth, river bed and flow speed).

This fieldwork suggests the water in Fohisokina has good physico-chemical qualities upstream according to Malagasy legislation drinking standards, but still needs microbiological analyzes to understand whether

it is potable. The source surface is still covered by plants and mosses, herbs and forest relicts. In contrast, the downstream water is progressively bad due to the degradation of the environment by bushfire and erosion during the rainy season; this increases the turbidity of the water and contamination.

Acknowledgements

We are grateful to the stakeholders who supported the project, among which are Conservation International, Malagasy Administration forest, Man and Environment, VOI FOMISAME, Rural Commune of Ivato, Mohamed bin Zayed Species Conservation Fund and Brother Industries.

References

1. N. Rabibisoa, *Mantella cowanii* Action Plan. Conservation International, Antananarivo, Madagascar (2008); available at www.sahonagasy.org
2. F. Andreone, F. C. E. Rabemananjara, C. Weldon. Policymaking and preventing the introduction of chytrid fungus in Madagascar. World Congress of Herpetology, Vancouver (2012). Abstract, 21.
3. N. Rabibisoa. Ny ela manova raha (Conservation International, Antananarivo, 2011).



Fig. 4: First carp production from Fohisokina fishpond.